



Geochemistry of Ediacaran-Ordovician diabase dikes in southern Colorado with a potential relationship to rifting in the Southern Oklahoma aulacogen

Caleb Perkey¹, Richard Hanson¹, Westin Scott², Benjamin Magnin³, Allen Stork⁴ and Yvette Kuiper⁵

¹Department of Geological Sciences, Texas Christian University, Fort Worth, TX 76129; ²Department of Geology, University of Kansas, Lawrence, KS 66045; ³Geology, Geophysics, and Geochemistry Science Center, U.S. Geological Survey, Denver Federal Center, Denver, CO 80225; ⁴Department of Natural and Environmental Sciences, Western Colorado University, Gunnison, CO 81231; ⁵Department of Geology and Geological Engineering, Colorado School of Mines, Golden, CO 80401



Abstract

The Southern Oklahoma aulacogen is a northwest-trending structure containing abundant igneous rocks representing the remains of a major Cambrian rift zone. Previous geologists have mapped numerous igneous intrusions in Colorado that follow the same trend, ranging from Ediacaran to Ordovician in age, and have speculated that these intrusions may be a part of the same rift. These intrusions include abundant igneous dikes of various compositions that originated from deeper magmatic bodies, filling fracture systems in older igneous rocks and Precambrian gneisses. This study involves the geochemical analysis of samples we collected from diabase dikes found along that northwest trend in southern Colorado. The dikes include a prominent diabase dike swarm in the Gunnison area as well as other individual dikes in the Wet Mountains and Front Range farther east. On the discrimination and REE diagrams, twenty-six representative dike samples from both sample regions plot tightly together, indicating the clustered dikes share a petrogenetic history of E-MORB-type magma that interacted with intercontinental lithosphere. In addition, this cluster generally plots within the same regions as data from diabase dikes associated with the Southern Oklahoma aulacogen.

Fifteen samples were taken from generally NW-trending diabase dikes in the Gunnison dike swarm, and these make up the majority of the dike samples that cluster together. The remaining eleven samples originate from general NW-trending diabase dikes in other locations across southern Colorado. Five samples were taken from WNW- to NW-trending diabase dikes in the Wet Mountains. Five samples were taken from NW- to NNW-trending diabase dikes in the Front Range, ~80 km north of the Wet Mountains. One sample was taken from a NW-trending diabase dike in the Unaweep Canyon, ~270 km west-northwest of the Wet Mountains. In addition, a sample was taken from a Cenozoic-age basalt dike for comparison with the samples from diabase dikes.

The geochemical similarities between diabase dikes sampled for this study and those within the Southern Oklahoma aulacogen suggest a linked petrogenetic history. Furthermore, the distribution of these samples raises the intriguing possibility that dikes related to Ediacaran-Ordovician intraplate magmatism in Colorado may be more extensive than previously thought.

Geologic Background

Cambrian igneous rocks in southern Oklahoma and adjacent parts of Texas, mostly present in the subsurface, define a northwest-trending rift zone extending from the ancient continental margin (Hanson et al., 2013; Wall et al., 2021). Numerous Ediacaran to Ordovician igneous intrusions ranging from large plutonic intrusions to smaller dikes occur along the same trend in Colorado, primarily in the Powderhorn District in western Colorado and the Wet Mountains and other parts of the Front Range of the Rockies farther east (Figs. 1 and 2). It is possible that all these rocks form a single major rift, where North America began to split apart (Larson et al., 1985). This project deals with abundant dikes present in the Gunnison region and the Wet Mountains and parts of the Front Range farther east (Olson et al., 1977; Pivarunas and Meert, 2019; Magnin et al., 2023). We have sampled a range of dikes mapped by previous workers that consist of diabase, lamprophyre and what was thought to be trachyte (see unnumbered field photos to lower left and upper right), as well as one much younger basaltic dike, in order to compare it with the older dikes. Associated plutons (Fig. 2) include the Powderhorn Complex, which contains dolomitic carbonatite, nepheline syenite, and other alkalic rocks. The McClure Mountain Complex in the Wet Mountains consists of a layered mafic-ultramafic complex, syenite, nepheline syenite, and a pyroxene-nepheline rock.




Fig. 1 Map of Colorado indicating the locations of igneous complexes and dikes thought to be associated with the Southern Oklahoma aulacogen. Modified from Larson et al. (1985).



Caption



Caption

Methods and Sampled Units

We collected samples in Colorado (Fig. 3) during the summers of 2022, 2023, and 2025. These samples are from diabase dikes in the Royal Gorge, Front Range, and Gunnison area, as well as alkaline and additional diabase dikes scattered throughout the Wet Mountains region. We also collected a Cenozoic basalt dike in the Wet Mountains for purposes of comparison with our other samples. Samples were prepared at TCU and then sent to the GeoAnalytical Lab at Washington State University. Major and trace elements were analyzed by X-ray fluorescence spectrometry and inductively coupled plasma-emission mass spectrometry.




Fig. 5 Map of the Front Range region with arrows indicating where diabase samples were collected. Modified from Ruleman et al. (2011).




Fig. 4 Map of the Royal Gorge region with arrows indicating where diabase samples were collected. Modified from Taylor et al. (1975).




Fig. 3 Geologic Map of Colorado, United States (Tweet, 1979).




Fig. 6 Map of the Gunnison area of western Colorado with sample locations (Stork, 2022, unpublished data).

Geochemistry

Winchester and Floyd Classification Diagram

We used the Winchester and Floyd (1977) classification diagram for igneous rocks (Fig. 4) because it uses ratios of highly immobile elements that tend to resist secondary alteration. Most of the diabase samples fall within the subalkaline basalt field. However, three diabase samples fall within alkaline fields. Two of these samples fall within the alkaline basalt field, and another sample falls within the basanite/nephelinite field. Another sample (23-12) plots far to the left of the sub-alkaline basalt field, which matches its erratic behavior in other diagrams. The lamprophyre dike samples plot within the alkaline basalt and basanite/nephelinite fields, as does the mafic-rich nepheline syenite dike. Samples from dikes that were previously classified as red trachytes by other workers plot within the phonolite field, suggesting these dikes were previously misidentified.




Fig. 5 Discrimination diagram for basalts, from Pearce (1982).




Fig. 6 Discrimination diagram for basalts, from Pearce and Cann (1973).




Fig. 7 Discrimination diagram for basalts, from Cabanis and Lecomte (1989).




Fig. 8 Discrimination diagram for basalts, from Meschede (1986).




Fig. 9 Discrimination diagram from Wood (1980).




Fig. 10 Discrimination diagram from Pearce (2008).

REE (Rare-Earth Element) Diagrams

The majority of the Gunnison area and Wet Mountains samples cluster closely together with moderate light REE-enriched patterns in the upper graph of Fig. 11, typical of E-MORB-type basalts. The enrichment typical of these alkaline magmas. The anomalous sample (23-12) shows an unusual REE pattern that we do not understand. In the lower graph of Fig. 11, the phonolite dike samples show a well-defined europium anomaly along with a flat heavy REE pattern. This pattern contrasts with the trends of the lamprophyre and mafic-rich nepheline syenite dike samples, all of which show a much steeper light REE-enriched pattern. In Fig. 12, the diabase dike samples from the Gunnison area and the Wet Mountains/Front Range mostly show similar patterns. The phonolite dike samples show strong negative anomalies in barium, strontium, and titanium, suggesting these dikes are the product of prolonged fractional crystallization. The phonolite dike samples also show enrichment of many of the incompatible elements on the left side of the diagram, which is typical of alkaline rocks. The diabase sample (23-12) is completely anomalous.




Fig. 11 REE diagrams of dike samples normalized to primitive mantle (Sun and McDonough, 1989).

Discrimination Plots

Various discrimination plots employing immobile trace elements show most of the diabase samples from the Gunnison area and the Wet Mountains/Front Range falling within the fields for tholeiitic, within-plate, continental basalts (Figs. 5-8). Samples that plot within the alkaline fields in the Winchester and Floyd (1977) diagram also tend to fall within alkaline fields in Fig. 8. In Fig. 9, most of the diabase samples have E-MORB-type affinities. This is consistent with the results shown in Fig. 10, which indicates the magmas are E-MORB-type, asthenospheric partial melts that interacted with continental lithosphere previously modified by subduction. Three alkaline basalt samples have OIB-type affinities. The anomalous sample (23-12) continues to behave erratically.




Fig. 12 REE diagram of all dike samples normalized to primitive mantle (Sun and McDonough, 1989).

Conclusions

- The samples fall within the sub-alkaline basalt, alkaline basalt, basanite/nephelinite, and phonolite fields in Fig. 4. This diagram indicates that the dike samples previously classified as trachyte dikes by other workers are, in fact, phonolites.
- All these rock types are typical of intraplate rift environments. Figs. 5-8 show that the majority of the diabase dikes plot within fields for tholeiitic, within-plate, continental basaltic magmas. Figs. 9 and 10 show that the parental magmas of the dikes were E-MORB-type, asthenospheric melts that interacted with continental lithosphere previously modified by subduction.
- The lack of any observable trends in the Harker variation diagrams (Fig. 3) at least partly reflects the fact that we are plotting samples that have different petrogenetic histories as well as the fact that the samples have been altered to various degrees. One diabase sample (23-12) shows erratic and anomalous behavior across the various geochemical diagrams, and this must also reflect secondary alteration, although the cause of that alteration is unclear.
- Figs. 11 and 12 show that most of the diabase samples exhibit E-MORB-type patterns, and the alkaline basalts show light REE enrichment. However, the anomalous diabase sample (23-12) shows a very unusual REE pattern. The phonolite dike samples all show well-defined, negative europium anomalies as well as negative anomalies in barium, strontium, titanium, indicating the parental magmas underwent prolonged fractional crystallization. The lamprophyre and mafic-rich nepheline syenite dike samples show very similar REE patterns with pronounced light REE enrichment and appear to be unrelated to the phonolites.
- The compositions of the Colorado dikes conflict with the dominant rock types in the Southern Oklahoma aulacogen, which include large volumes of rhyolite and granite. The diabase dikes with E-MORB-type affinities in Colorado are geochemically very similar to argues in the Southern Oklahoma aulacogen (e.g., Eschberger et al., 2014), which could argue for a direct connection between the two igneous suites. However, lamprophyres, phonolites, and nepheline syenites do not occur in the aulacogen. If the aulacogen does directly continue into Colorado, then it must have had a different petrogenetic history. Isotopic dating is now in progress to determine if there is a direct connection between the two regions.

References

Armbrustmacher, T. J., 1979. Replacement and Primary Magmatic Carbonates from the Wet Mountains Area, Fremont and Custer Counties, Colorado. *Economic Geology*, v. 74, p. 888-901.

Cabanis, B., Lecomte, M., 1989. Le diagramme La³/Y³-Nb⁵/B: un outil pour la discrimination des séries volcaniques et la mise en évidence des processus de mélange et/ou de contamination crustale. *Comptes Rendus de l'Académie des Sciences, Série 2, Mécanique, Physique, Chimie, Sciences de l'Univers, Sciences de la Terre*, v. 309, p. 2023-2029.

Hanson, R., Puckett, R., Keller, C., Brueske, M., Bolen, C., Martzman, S., Finagan, S., McCleary, D. A., 2013. Intraplate magmatism related to opening of the southern Iapetus Ocean: Cambrian Wichita igneous province in the Southern Oklahoma rift zone. *Lithos*, v. 174, p. 57-70.

Eschberger, A. M., Hanson, R. E., Puckett, R. E., 2014. Carlton Rhyolite Group and diabase intrusions in the East Timberr Hills, Arbuckle Mountains, in Suneson, N. (ed.), *Igneous and Tectonic History of the Southern Oklahoma Aulacogen*. Oklahoma Geological Survey Guidebook 38, p. 143-186.

Larson, E. E., Patterson, P. E., Curtis, G., Drake, R., Mutschler, F. E., 1985. Petrologic, paleomagnetic, and structural evidence of a Paleozoic rift system in Oklahoma, New Mexico, Colorado, and Utah. *Geological Society of America Bulletin*, v. 96, p. 1364-1372.

Magnin, B. P., Kuiper, Y. D., Anderson, E., 2023. Ediacaran-Ordovician magmatism and REE mineralization in the Wet Mountains, Colorado, USA: implications for failed continental rifting. *Tectonics*, Vol. 42, Article e2022TC007674.

Meschede, M., 1986. A method of discriminating between different types of mid-ocean basalts and continental tholeiites with the Nb-Zr-Y diagram. *Chemical Geology*, v. 56, p. 297-318.

Olson, J. C., Marvin, R. F., Parker, R. L., Mehnert, H. H., 1977. Age and tectonic setting of lower Paleozoic alkalic and mafic rocks, carbonatites, and thorium veins in south-central Colorado. *Journal of Research of the U. S. Geological Survey*, v. 5, p. 673-687.

Pearce, J. A., Cann, J. R., 1973. Tectonic setting of basic volcanic rocks determined using trace element analysis. *Earth and Planetary Science Letters*, v. 19, p. 290-300.

Pearce, J. A., 1982. Trace element characteristics of lavas from destructive plate margins: Andesites; Orogenic andesites and related rocks, p. 525-548.

Pearce, J. A., 2008. Geochemical fingerprinting of oceanic basalts with applications to ophiolite classification and the search for Archean oceanic crust. *Lithos*, v. 100, p. 14-48.

Pivarunas, A. F., Meert, J., 2019. Protracted magmatism and magnetization around the McClure Mountain alkaline igneous complex: Lithosphere, v. 11, p. 590-620.

Ruleman, C. A., Bohannon, R. G., Bryant, B., Shroba, R. R., Premo, W. R., 2011. Geologic Map of the Bailey 30' x 60' Quadrangle, North-Central Colorado: USGS Scientific Investigations Map 3155.

Sun, S. S., McDonough, W. F., 1989. Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. *Geological Society Special Publications*, v. 42, p. 313-345.

Wall, C., Hanson, R., Schmitz, M., Price, J., Donovan, R., Boro, J., Eschberger, A., Toews, C., 2021. Integrating zircon trace-element geochemistry and high-precision U-Pb zircon geochronology to resolve the timing and petrogenesis of the late Ediacaran-Cambrian Wichita igneous province, Southern Oklahoma aulacogen, USA. *Geology*, v. 49, p. 268-272.

Winchester, J. A., Floyd, P. A., 1977. Geochemical discrimination of different magma series and their differentiation products using immobile elements. *Chemical Geology*, v. 20, p. 325-343.

Wood, D. A., 1980. The application of a Th-Hf-Ta diagram to problems of tectonomagmatic classification and to establishing the nature of crustal contamination of basaltic lavas of the British Territory volcanic province. *Earth and Planetary Science Letters*, v. 50, p. 11-30.